

## 2. Biology (from Engineering Perspective)

Read:

Chapter 2  
(pp. 10 - 55)

# A. Cell types

## 1. Viruses

- obligate parasites
- 30 - 200 x 10<sup>-9</sup> m diameter (0.03 - 0.2 μm)
- cause infections in other higher cells
- **Bacteriophages** are viruses which specifically infect bacteria.
- contain DNA or RNA enclosed in a protein coat called a **capsid**

- After infection:
  - 1) **Lytic Cycle** - host cells lyse (break apart) and release new viruses, which go on to infect new host cells.
  - 2) **Lysogenic Cycle** - viral DNA is incorporated into host DNA, and continues to multiply with host.
- Viruses are important because they can:
  - 1) infect bacteria used in a commercial process.
  - 2) cause disease in animals and plants.
  - 3) be used to incorporate foreign DNA into a host cell.

## 2. Prokaryotes

- 5 - 30 x 10<sup>-7</sup> m diameter (0.5 - 3 μm)
- do not contain a membrane-enclosed nucleus
- usually do not associate with other cells
- often multiply quickly when nutrients are plentiful  
Some populations can double every 20 minutes.

## a. Eubacteria

- categorized by their cell envelope

### Gram-negative

8 nm cytoplasmic membrane

7 nm periplasmic space

2 nm peptidoglycan

8 nm outer membrane

### Gram-positive

8 nm cytoplasmic membrane

80 nm peptidoglycan

Secretion of proteins is often more difficult for Gram-negative cells because of outer membrane.

Gram stain developed to distinguish these cell types, involves a dye which is retained by peptidoglycan.

- formation of spores in response to adverse conditions
- formation of inclusion bodies (can be large)  
examples: sulfur, proteins
- excretion of polysaccharides to adhere to surfaces  
(teeth, bioremediation)

#### b. Archaeobacteria

- have no peptidoglycan
- have different lipid composition in cytoplasmic membrane
- live in extreme environments  
examples: 110°C, dead sea water

### 3. Eukaryotes

- 1 - 20 x 10<sup>-6</sup> m diameter (1 - 20 μm)
- contain a membrane-enclosed nucleus
- Cell membrane has sterols which provide rigidity.
- Plant cells contain cellulose fibers which make them strong. Animal cells do not have cellulose.
- have numerous complex cell organelles
  - 1) **Mitochondria** are site for respiration and oxidative phosphorylation. **Cristae** are their complex inner membrane folds.  
Have their own DNA  
Replicate independently
  - 2) **Chloroplasts** are site for photosynthesis.

Both eukaryotes and prokaryotes contain:

### Endoplasmic reticulum

- **rough** - Contain ribosomes and are site of protein synthesis
- **smooth** - site of lipid synthesis


Both eukaryotes and prokaryotes may contain:

**Flagella** and **cilia** are filamentous structures which permit cell to propel through fluid. **Chemotaxis** is the tendency of cells to move towards/away from chemicals.

## a. Fungi


### 1) Yeasts

- spherical cells
- widespread in nature (soil)
- do not use sunlight for energy
- free-living
- reproduce by **budding**




## 2) Molds

- distinctly non-spherical cells
- A **Mycelium** is a highly branched system of hyphae.
- **Hyphae** are thin filaments of cells.
- **Conidia** are asexual spores which form on hyphae.
- in submerged culture can form thick mats which result in complex rheological behavior




Penicillium



Septate Hypha


Rhizoctania



Conidiophore

Conidia

Aspergillus



Chrysosporium

## 2) Molds

- distinctly non-spherical cells
- **Mycelium** are a highly branched system of tubes.
- **Hyphae** are thin filaments of cells.
- **Conidia** are asexual spores which form on hyphae.
- in submerged culture can form thick mats which result in complex rheological behavior
- have high oxygen requirement
- do not use sunlight for energy
- free-living
- spores

## b. Algae

- photosynthetic
- often, but not always, unicellular
- some contain silica or calcium carbonate in structure
- principal products:
  - diatomaceous earth
  - agar
  - foodstuffs


### c. Protozoa

- large unicellular cells that lack cell walls
- ingest other smaller organisms
- move to seek prey
  - amebae** - cytoplasm forms pseudopodium
  - flagellates** - have one flagellum
  - ciliates** - have many cilia
  - sporozoa** - usually have aid of host

### d. Animal and Plant Cells

- used for many vaccines (animal cells)
- therapeutic proteins (animal cells)
- colors, flavors, insecticides (plant cells)

## Comparison of cell growth rates




## B. Cell construction

### 1. Proteins


- composed of 20 amino acids covalently linked
- have diverse functions:

<u>Class</u>	<u>Example</u>	<u>Function</u>
enzymes	alcohol dehydrogenase	oxidizes alcohols to aldehydes
regulatory	insulin	regulates glucose metabolism
transport	hemoglobin	transport of oxygen and carbon dioxide in blood
protection	IgG antibody	forms complex with foreign protein and initiates immune response
storage	albumin	egg-white
structure	collagen	cartilage and tendons


## 2. Amino acids




- **zwitterion** - doubly charged
- chiral - L-form found in proteins
- **Isoelectric point** is pH at which amino acid has no net charge.
- **R** = side chain  
The R-group distinguishes amino acids.
- Proteins are formed from amino acids by peptide bonds



Glycine




Alanine




Isoleucine



Aspartate



Phenylalanine



Lysine

### 3. Protein structure


- **Primary structure** is merely the sequence of amino acids
- **Secondary structure** is the structural conformation of amino acids that are close neighbors. The peptide bond restricts flexibility.
- **Tertiary structure** is the three dimensional structure arising from hydrogen bonds, disulfide bridges, hydrophobic and ionic interactions between amino acids which are widely separated in linear sequence but are spatially proximate.
- **Quarternary structure** is the interactions between protein subunits.

## 4. Carbohydrates

- General formula is  $(\text{CH}_2\text{O})_N$ .
- Involved in structure, storage and signaling
- Analogous to amino acids, monomeric units can combined to form polymeric molecules called **polysaccharides**


## a. Monosaccharides

- 3-9 carbon atoms
- 5 or 6 carbon atoms most common, which often form stable **furanose** (5-membered) or **pyranose** (6-membered) ring
- examples:




$\beta$ -D-fructose




Haworth  
projection




$\alpha$ -D-glucose




$\beta$ -D-glucose




$\alpha$ -D-glucose  
 $\alpha$  = below ring



$\beta$ -D-glucose  
 $\beta$  = above ring



$\alpha$ -D-glucopyranose



$\beta$ -D-glucopyranose

35%  $\alpha$   
64%  $\beta$   
1% furanose  
<0.03% linear


## b. Disaccharides

- formed by the condensation of two monosaccharides


(while amino acids only form one type of covalent bond - peptide bond - saccharides can form several different linkages)


- examples:

$\alpha$ -1,4 linkage  
maltose




$\beta$ -1,4 linkage  
cellobiose






Maltose



Cellobiose




$\beta$ -D-galactose

+

$\beta$ -D-glucose

with  $\beta$ -1,4 linkage yields lactose

$\beta$  (1 $\rightarrow$ 4) galacto-glucopyranose




$\alpha$ -D-glucose

+


$\beta$ -D-fructose

with  $\alpha, \beta$ -1,2 linkage yields sucrose

$\alpha, \beta$  (1 $\rightarrow$ 2) gluco-fructofuranose



Lactose



Sucrose

### c. Polysaccharides

- formed by the condensation of more than two monosaccharides
- examples:

Amylose      Straight chain of 15 - 3,000 glucose units linked by  $\alpha$ -1,4 glycosidic bonds.

**Insoluble in water**

Cellulose      Straight chain of 300 - 6,000 glucose units linked by  $\beta$ -1,4 glycosidic bonds.

**Insoluble in water**

### c. Polysaccharides (cont'd)

Amylopectin      Branched chain of 5,000 - 10,000 glucose units linked by  $\alpha$ -1,4 and  $\alpha$ -1,6 glycosidic bonds. Branching occurs via  $\alpha$ -1,6 bonds after an average of 25 units.

More exposed monomers available for enzymatic cleavage.


**Soluble in water (forms gels)**

Glycogen      Like amylopectin but smaller and more frequent branching.


### c. Polysaccharides (cont'd)

Starch	20% amylose 80% amylopectin
--------	--------------------------------


Cellulose is more difficult to decompose in nature than amylose and amylopectin because of *interchain* hydrogen bond between C3 hydroxyl and pyranose oxygen leading to **crystalline** structure.



amylose



cellulose



amylopectin

## c. Polysaccharides (cont'd)

### Production of Sweeteners




### Relative Sweetness

glucose	0.7
sucrose	1.0
fructose	2.0

How to obtain more fructose?

### c. Polysaccharides (cont'd)



## 5. DNA

- long threadlike molecule which carries genetic information in its sequence of **purine** and **pyrimidine** bases
  - **transcription:** DNA template to messenger RNA
  - codon on mRNA interacts with anticodon on transfer RNA
  - **translation:** RNA code converted to amino acid sequence

## 5. DNA (cont'd)

- **plasmids** are nonchromosomal self-replicating segments of DNA floating around in cytoplasm
  - plasmids are a means for cells to transfer DNA quickly between cells and between species (e.g., give resistance to cells against antibiotics).
  - plasmids can be inserted into cells to get them to overproduce a desired protein.
  - **copy number** is the number of desired plasmids in a cell.

## C. Cell Nutritional Requirements

### 1. Energy

- phototrophs utilize light energy
- chemotrophs utilize chemical energy
  - lithotrophs utilize inorganic chemicals
  - organotrophs utilize organic chemicals

### 2. Carbon

- autotrophs utilize inorganic carbon ( $\text{CO}_2$ )
- heterotrophs utilize organic carbon

Industrial fermentations use cheap carbon sources such as molasses, malt extract, corn steep liquor, sulfite waste liquor

### “complex” media

Nature of carbon source has great impact on growth of cells:

### Example (with *E. coli*)

*J. Bacteriol.* (1971) 105:20+

complex media

Doubling time

35 min

glucose + salts

50 min

acetate + salts

80 min

succinate + salts

300 min

} “defined”  
media

## Advantages of using complex media instead of defined media:

- 1) Cheap
- 2) Readily available
- 3) Best cell growth
- 4) Have all nutritional requirements

## Disadvantages of using complex media instead of defined media:

- 1) Considerable variation in media composition
- 2) Processes less reliable and reproducible
- 3) Complicates separation and purification

### 3. Nitrogen

12% of bacterial dry mass

7.5% of yeast dry mass

- **complex media**
  - yeast extract  
(Baker's yeast lysed at 55°C)
  - peptone (protein hydrolysate)  
meat, cottonseeds, soy
- **defined media**
  - ammonia
  - ammonium salts
  - urea

## Prices for Defined Nitrogen Sources (Dec 2002)

<u>Substance</u>	<u>%N</u>	<u>\$ per lb</u>	<u>\$ per lb N</u>
NH <sub>3</sub>	0.823	0.073	0.088
NH <sub>4</sub> Cl	0.262	0.21	0.802
(NH <sub>4</sub> ) <sub>2</sub> SO <sub>4</sub>	0.212	0.053	0.250
NH <sub>4</sub> H <sub>2</sub> PO <sub>4</sub>	0.122	0.082	0.672 1)
(NH <sub>4</sub> ) <sub>2</sub> HPO <sub>4</sub>	0.212	0.062	0.292 2)
NH <sub>4</sub> NO <sub>3</sub>	0.350	0.069	0.197
urea	0.46	0.054	0.117

1) 27% P - \$0.30/lb P

2) 24% P - \$0.26/lb P

Are materials pure enough for biological processes?

## 4. Phosphorus

3% of bacterial dry mass

1.5% of yeast dry mass

- phosphate salts added if more P is necessary
  - 25% as much P is needed as N
  - 180% as much P is supplied as N
- in some fermentations, P can inhibit production of desired product

## 5. Sulfur

1% of bacterial and yeast dry mass

- sulfate salts added if more S is necessary
  - 10% as much S is needed as N
  - 60% as much S is supplied as N

## 6. Micronutrients

### a) trace elements:

$K^+$ ,  $Mg^{++}$ ,  $Ca^{++}$ ,  $Fe^{++}$ ,  $Mn^{++}$ ,  $Co^{++}$ , etc.

- supplied as necessary (in excess?) by addition of salts
- very important in regulating enzymes (both presence and absence)
- carefully controlled (example: citric acid fermentation)

### b) vitamins

- expensive

Elemental Composition of *E. coli*  
(mg / g dry cell weight)

<u>Element</u>		<u>growth</u>		<u>osmol,</u>	
		<u>early</u>	<u>late</u>	<u>low</u>	<u>high</u>
C	---	393	---	---	---
N	---	140	---	---	---
P	---	41	55	---	---
S	---	10	13	---	---
Mg	2.8	6.4	8.7	---	---
K	12	41	22	18	40
Na	---	23	53	3.9	3.2
ref	(1)	(2)		(3)	

Elemental Composition of *E. coli*  
(mg / g dry cell weight)

<u>Element</u>		<u>growth</u>	
		<u>early</u>	<u>late</u>
Ca	2.3	2.1	0.5
Cl	---	23	25
Zn	0.15	41	55
ref	(1)	(2)	

(1) Kung et al. *J. Bacteriol.* 1976, 126(3):1089

(2) Heldal et al. *Appl. Environ. Microbiol.* 1985 50(5):1251

(3) Epstein & Schultz *J. Gen. Physiol.* 1965 49:221

## Comments

- composition of cells is very flexible...cells have ability to adapt to environment.
- composition of medium can strongly impact composition of cells, which in turn can profoundly impact cellular metabolism (and product formation).
- ~20-30% of carbon consumed by an organoheterotroph ends up as cell material. Thus, 3-5 times as much carbon needs to be added as calculated merely from cell composition.